

Quasiparticle damping in two-dimensional superconductors with unconventional pairing

M. L. Titov, A. G. Yashenkin and D. N. Aristov

Petersburg Nuclear Physics Institute, Gatchina, St. Petersburg 188350, Russia.

(February 1, 2008)

Abstract

We calculate the damping of excitations due to four-fermionic interaction in the case of two-dimensional superconductor with nodes in the spectrum. At zero temperature and low frequencies it reveals gapless ω^3 behavior at the nodal points. With the frequency increasing the crossover to the normal-state regimes appears. At high frequencies the damping strongly depends on details of a normal-state spectrum parametrization. Two important particular cases such as the models of almost free and tight-binding electrons are studied explicitly and the characteristic scales are expressed through the model-free parameters of the spectrum at the nodal points. The possibility of crossover in temperature dependence of damping in the superconducting phase is discussed.

I. INTRODUCTION

An assumption that the pairing state in the high-temperature superconductors is the d -wave-like one is widely discussed nowadays. [1] Although up to now it apparently cannot be treated as the uniquely established fact there exist a large body of experimental data which supports it. [2] Such a pairing would result in presence of “nodal” points in the superconducting excitations spectrum in whose vicinities it has gapless phonon-like character. It should essentially modify the damping behavior comparing to the s -wave pairing state.

The quasiparticle damping owing to impurities has been extensively studied in literature. [3] However, this is not the only channel, though, maybe, the more interesting one. The excitations scattering by phonons and also by each others always contribute to the damping, as well.

In the present paper we evaluate the quasiparticle damping in two-dimensional (2D) superconductors with d -wave-like pairing appeared due to their four-fermionic interaction. It may be the on-site Coulomb repulsion in the small- U Hubbard model, or the residual part of the interaction which provides the pairing in the strong correlations scheme. [4,5]

In the BCS case, the damping at zero temperature has the threshold at $\omega = 3\Delta$. We show that for the gap function with nodes it demonstrates gapless ω^3 low-frequency behavior at the nodal points. The results obtained for high frequencies depend on details of Fermi surface (FS) parametrization. The damping for the circular FS will increase quadratically while for the spectrum with Van-Hove singularities the additional linear regime appears. Being generalized onto finite temperatures, this results allow us to conclude about existence of crossovers in temperature dependence of damping in the superconducting state.

The outline of this paper is as follows. In Sect. II we obtain general expression for damping of superconducting excitations due to their interaction. In Sect. III we evaluate it for the case of 2D superconductor with nodes in the spectrum. Since the high-frequency behavior of damping depends strongly on FS parametrization, in the consequent subsections we analyze three principal cases: (i) the case of circular FS and also the case of FS of varying

curvature when the gap node intersects it (ii) at the almost flat piece or (iii) in the vicinity of saddle-point of the spectrum. At last, in Sect. IV we generalize and discuss the results obtained.

II. GENERAL FORMALISM

It is convenient to represent the Hamiltonian $\mathcal{H} = \mathcal{H}_o + \mathcal{H}_i$ in the Nambu notation

$$\begin{aligned}\mathcal{H}_o &= \sum_{\mathbf{k}} \Psi_{\mathbf{k}}^+ (\xi_{\mathbf{k}} \tau_3 + \Delta_{\mathbf{k}} \tau^- + \Delta_{\mathbf{k}}^* \tau^+) \Psi_{\mathbf{k}}, \\ \mathcal{H}_i &= \frac{1}{2} \sum_{\mathbf{k}, \mathbf{k}', \mathbf{q}} V_{\mathbf{k}+\mathbf{q}, \mathbf{k}'-\mathbf{q}; \mathbf{k}, \mathbf{k}'} (\Psi_{\mathbf{k}+\mathbf{q}}^+ \tau_3 \Psi_{\mathbf{k}}) (\Psi_{\mathbf{k}'-\mathbf{q}}^+ \tau_3 \Psi_{\mathbf{k}'}).\end{aligned}\tag{1}$$

Here $V_{\mathbf{k}_1 \mathbf{k}_2; \mathbf{k}_3 \mathbf{k}_4}$ is the interaction, $\Psi_{\mathbf{k}}^+ = (c_{\mathbf{k}\uparrow}^\dagger, c_{-\mathbf{k}\downarrow})$ stands for the spinor with $c_{\mathbf{k}\sigma}^\dagger$ being the fermionic creation operator, $\xi_{\mathbf{k}}$ and $\Delta_{\mathbf{k}}$ are the normal-state conduction band and the gap function (GF), respectively, and $\tau^\pm = \frac{1}{2}(\tau_1 \pm i\tau_2)$, where τ_i denote the Pauli matrices.

Let us define the bare matrix Green function as follows

$$\hat{G}^{(0)} = \begin{pmatrix} G^{(0)} & -F^{(0)} \\ F^{(0)+} & G^{(0)+} \end{pmatrix}\tag{2}$$

where normal G and anomalous F Green functions are written as

$$G_{\mathbf{k}}^{(0)}(i\omega_n) = -\frac{i\omega_n + \xi_{\mathbf{k}}}{\omega_n^2 + \varepsilon_{\mathbf{k}}^2}, \quad F_{\mathbf{k}}^{(0)}(i\omega_n) = \frac{\Delta_{\mathbf{k}}}{\omega_n^2 + \varepsilon_{\mathbf{k}}^2}.\tag{3}$$

Here $i\omega_n = 2\pi i(n + \frac{1}{2})T$ is the imaginary Matsubara frequency and the excitation spectrum is given by $\varepsilon_{\mathbf{k}} = [\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2]^{1/2}$.

Within this notation the Dyson equation obtains the form $\hat{G}^{-1} = \hat{G}^{(0)-1} - \hat{\Sigma}$, with the diagonal self-energy operator $\hat{\Sigma} = \text{diag}(\Sigma, \Sigma^+)$.

The first nonvanishing contribution to Σ is given by

$$\begin{aligned}\Sigma(\mathbf{k}, i\omega_n) &= T^2 \sum_{\mathbf{1}, \mathbf{2}, \mathbf{3}} V_{1,2;3,0} V_{3,0;1,2} \delta_{1,2;3,0} \\ &\times \left[G_{\mathbf{1}}^{(0)} G_{\mathbf{2}}^{(0)} G_{\mathbf{3}}^{(0)} + F_{\mathbf{1}}^{(0)} F_{\mathbf{2}}^{(0)+} G_{\mathbf{3}}^{(0)+} \right],\end{aligned}\tag{4}$$

where the indices **1, 2, 3** label the intermediate frequencies and momenta and **0** corresponds to the external one.

After the analytic continuation $i\omega_n \rightarrow \omega + i\delta$ for the imaginary part of the self-energy at $T = 0$ and $\omega > 0$ we get

$$\begin{aligned} \text{Im}\Sigma(\mathbf{k}, \omega) = & -\pi \sum_{\mathbf{q}, \mathbf{p}} \int_0^\omega dx \int_0^{\omega-x} dy V_{\mathbf{k}, \mathbf{q}+\mathbf{p}-\mathbf{k}; \mathbf{q}, \mathbf{p}} V_{\mathbf{q}, \mathbf{p}; \mathbf{k}, \mathbf{q}+\mathbf{p}-\mathbf{k}} \\ & \times \delta(x - \varepsilon_{\mathbf{q}}) \delta(y - \varepsilon_{\mathbf{p}}) \delta(x + y - \omega + \varepsilon_{\mathbf{q}+\mathbf{p}-\mathbf{k}}) \\ & \times [u_{\mathbf{q}}^2 u_{\mathbf{p}}^2 v_{\mathbf{q}+\mathbf{p}-\mathbf{k}}^2 - u_{\mathbf{q}} u_{\mathbf{p}}^* v_{\mathbf{q}} v_{\mathbf{p}}^* u_{\mathbf{q}+\mathbf{p}-\mathbf{k}}^2], \end{aligned} \quad (5)$$

where the coefficients of Bogolyubov transform are defined by

$$u_{\mathbf{q}}^2 = \frac{1}{2} \left(1 + \frac{\xi_{\mathbf{q}}}{\varepsilon_{\mathbf{q}}} \right), \quad v_{\mathbf{q}}^2 = \frac{1}{2} \left(1 - \frac{\xi_{\mathbf{q}}}{\varepsilon_{\mathbf{q}}} \right). \quad (6)$$

III. CALCULATIONS OF DAMPING

In this section we will evaluate Eq. (5) in the case of 2D superconductor with unconventional pairing assuming different model parametrizations for the Fermi surface.

We shall suppose that the gap function has nodal lines $|\Delta_{\mathbf{q}}| = 0$ in the 2D Brillouin zone. Let these lines intersect Fermi surface at a set of the points $\mathbf{q} = \mathbf{q}_n$. Then in the vicinities of such points one can write

$$\begin{aligned} \xi_{\mathbf{q}} &\simeq \mathbf{v}_F \bar{\mathbf{q}}, & \Delta_{\mathbf{q}} &\simeq \mathbf{v}_g \bar{\mathbf{q}}; \\ \mathbf{v}_F &= \left. \frac{\partial \xi_{\mathbf{q}}}{\partial \bar{\mathbf{q}}} \right|_{\bar{\mathbf{q}}=0}, & \mathbf{v}_g &= \left. \frac{\partial \Delta_{\mathbf{q}}}{\partial \bar{\mathbf{q}}} \right|_{\bar{\mathbf{q}}=0}, \end{aligned} \quad (7)$$

where $\bar{\mathbf{q}} = \mathbf{q} - \mathbf{q}_n$ and the index n labels the nodal point. We see that the excitation spectrum near these points has gapless linear character.

In general, the quasiparticle damping due to their interaction has to reveal threshold feature in frequency in the superconducting state. The threshold corresponds to the minimal energy of the excitation needed to decay into three quasiparticles in the intermediate state.

For the gap function with nodes however it disappears at the nodal points of the spectrum. Consequently, the damping at this points appear to be gapless. Taking it into account, we restrict ourselves in treatment of damping in the vicinities of nodal points only.

The straightforward calculations at $\omega \rightarrow 0$ yield

$$\text{Im } \Sigma(\mathbf{k}_n, \omega) \simeq -\lambda_n \omega^3$$

$$\lambda_n = \sum_{l,k} \frac{V_{n;lk}^2 \mathcal{F}_{lk}}{|\mathbf{v}_F^l \times \mathbf{v}_g^l| |\mathbf{v}_F^k \times \mathbf{v}_g^k|} \quad (8)$$

$$V_{n;lk}^2 = \lim_{\mathbf{k}, \mathbf{p}, \mathbf{q} \rightarrow \mathbf{k}_n, \mathbf{p}_l, \mathbf{q}_k} V_{\mathbf{k}, \mathbf{q} + \mathbf{p} - \mathbf{k}; \mathbf{q}, \mathbf{p}} V_{\mathbf{q}, \mathbf{p}; \mathbf{k}, \mathbf{q} + \mathbf{p} - \mathbf{k}},$$

where the indices l, k run over all the nodal points. Here $\mathcal{F}_{lk}$ are some dimensionless positive factors. The diagonal terms of the matrix $\hat{\mathcal{F}}$ (i.e., referring to a single nodal point) are always numbers of order of unity while the off-diagonal ones could have additional smallness for particular FS parametrizations via their dependence on spectrum parameters. The principal contribution to the damping at low frequencies comes from excitations with momenta lying in the small elliptic vicinities of nodal points (see Fig. 1) wherein linear approximation for the spectrum works well.

Note that we obtain the universal cubic frequency dependence of the excitations damping at the nodal points without appealing to any fixed parametrization for spectrum. By the way, such dependence means the stability of nodes with respect to this kind of interaction.

At the same time, some features of the model may, in principle, lead to different power laws. In particular, the damping would increase with frequency slower than ω^3 if at least one of spectrum zeroes (i) appeared due to tangency of FS and GF node, or (ii) coincided with van-Hove singularity in the band structure and/or (iii) with cusp- or discontinuity-point of the nodal line of the gap function. All these cases are obviously quite exotic.

When the frequency increases, the integration in Eq. (5) cannot be restricted to the nodal vicinities and such characteristics as general shape of FS, curvature of FS at the nodal points, existence of van-Hove singularities in the spectrum, *etc.*, should be taken into

account. One can distinguish three qualitatively different cases: (i) the circular FS and also the FS of varying curvature with the nodal points located (ii) far from or (iii) at the sharp curvature maximum. We shall illustrate the above cases by examples which would allow us to examine in detail the intermediate asymptotics and to express the corresponding crossover scales through the parameters of the models used.

A. Circular Fermi surface

Putting the lattice parameter to be unity, we write the circular FS as follows: $\xi_{\mathbf{q}} = q^2/2m - \varepsilon_F$. For definiteness, we choose the $d_{x^2-y^2}$ -wave gap function, $\Delta_{\mathbf{q}} = \Delta/2 [\cos q_x - \cos q_y]$. [5,6] In line with Eq. (8) we get:

$$\text{Im}\Sigma(\mathbf{k}_n, \omega) \simeq -\left(\frac{V}{\varepsilon_F}\right)^2 \frac{\omega^3}{\Delta^2}, \quad \omega \ll \frac{\Delta^2}{\varepsilon_F}, \quad (9)$$

where $V^2 \sim \max V_{n;ls}^2$. The principal contribution to $\text{Im}\Sigma$ is obtained when all the momenta $\mathbf{p}$, $\mathbf{q}$ and $\mathbf{k}_n$ are almost parallel and belong to the vicinities of the nodal points (see Fig. 1).

At higher frequencies $\Delta^2/\varepsilon_F \ll \omega \ll \varepsilon_F$ we find

$$\text{Im}\Sigma(\mathbf{k}_n, \omega) \simeq -\left(\frac{V}{\varepsilon_F}\right)^2 \frac{\omega^2}{\varepsilon_F} \ln \left[\min \left(\frac{\omega \varepsilon_F}{\Delta^2}; \frac{\varepsilon_F}{\omega} \right) \right] \quad (10)$$

To obtain Eq. (10), the curvature of FS was taken into account (see Appendix). The crossover from the cubic dependence to the quadratic one appears at $\omega \sim \Delta^2/\varepsilon_F$. It can be qualitatively explained on the following way. One can show that, when $\omega \ll \Delta^2/\varepsilon_F$, the areas of integration defined by arguments of δ -functions in Eq. (5) are located near the nodal points (see Fig. 1) where the spectrum can be well approximated by ellipses. When $\omega \sim \Delta^2/\varepsilon_F$, the areas of integration propagate along the whole Fermi surface so that not only the vicinities of the nodal points principally contribute to $\text{Im}\Sigma$. For $\Delta^2/\varepsilon_F \ll \omega \ll \Delta$, we obtain the quadratic intermediate asymptotic regime with unconventional logarithmic factor of the form $\ln(\omega \varepsilon_F/\Delta^2)$ which stems from integration over the FS. At frequencies higher then the gap value $\Delta \ll \omega \ll \varepsilon_F$ information about the superconducting state disappears,

and the conventional 2D Fermi-liquid result [7] is restored. Note that principal contributions to $\text{Im}\Sigma$ in the latter case again come from almost parallel $\mathbf{k}$, $\mathbf{p}$ and $\mathbf{q}$.

B. Tight-binding model, $d_{x^2-y^2}$ -wave pairing

Now we will demonstrate how the deformation of FS affects the damping. As before we shall consider $d_{x^2-y^2}$ gap function. Let us adopt the simplest tight-banding form for the normal-state dispersion: $\xi_{\mathbf{q}} = -2t [\cos q_x + \cos q_y] + \mu$. Here t is the nearest-neighbors hopping integral and μ is the chemical potential measured from the half-filling. Note that for small enough $\mu \ll t$ Fermi surface has long almost flat pieces near the points of intersection with the nodal lines of the $d_{x^2-y^2}$ gap function (see Fig. 2a).

Under the above conditions one can easily rewrite the low-frequency asymptotics of damping by substituting $\varepsilon_F \rightarrow t$ in the parenthesis and $\varepsilon_F \rightarrow \mu$ in other parts of Eq. (9). The range of validity for such expression is restricted by the condition $\omega \ll \min(\Delta^2/\mu, \Delta)$. Next, in order to obtain expression for damping at $\min(\Delta^2/\mu, \Delta) \ll \omega \ll \max(\mu, \Delta)$, the same substitutions in Eq. (10) should be done. Obviously, all the statements concerning the areas of principle contributions still hold. Finally, at $\max(\mu, \Delta) \ll \omega \ll t$ we obtain

$$\text{Im}\Sigma(\mathbf{k}_n, \omega) \simeq -\left(\frac{V}{t}\right)^2 \omega \quad (11)$$

Thus, the additional linear-in- ω high-frequency regime appears in this case. It occurs since at $\omega \gg \max(\mu, \Delta)$ the areas of integration in Eq. (5) engage the saddle-points of the normal-state spectrum located at $(\pm\pi, 0)$ and $(0, \pm\pi)$. For the normal state crossover to the linear behavior has been demonstrated in [8].

It follows from the aforementioned that there is no room for linear regime of damping at $\mu \sim t$ while the crossover from the cubic dependence to the quadratic one occurs at $\omega \sim \Delta^2/\mu$. This case can be treated as the one of slightly deformed circular FS without additional characteristic scales. When μ decreases (i.e., close to the half-filling) the long almost flat pieces of the FS appear as well as the saddle-points in the normal-state spectrum

come into play. For $\mu \ll \Delta \ll t$ there co-exist three ranges with cubic, quadratic and linear frequency dependencies matching at $\omega \sim \Delta^2/\mu$ and $\omega \sim \mu$, respectively. At last, for very small $\mu \ll \Delta$ linear approximation for spectrum works well unless the ellipses touch the saddle-points. Consequently, the range of quadratic dependence in the latter case disappears and there exist the only crossover scale $\omega \sim \Delta$.

C. Tight-binding model, d_{xy} -wave pairing

Finally, let us consider the situation when the nodal points lie at the points of maximum of FS curvature. To illustrate this case we write the normal-state spectrum in the tight-binding form and choose the gap function as d -wave with xy -symmetry, [9] $\Delta_{\mathbf{q}} = \Delta \sin q_x \sin q_y$ (see Fig. 2b). Then the straightforward calculations at $\omega \ll \mu$ yield the formulas similar to those obtained in previous subsection:

$$\text{Im}\Sigma(\mathbf{k}_n, \omega) \simeq -\left(\frac{V}{\mu}\right)^2 \frac{\omega^3}{\Delta^2}, \quad \omega \ll \frac{\Delta^2\mu}{t^2}. \quad (12)$$

When $\Delta^2\mu/t^2 \ll \omega \ll \mu$ we have

$$\text{Im}\Sigma(\mathbf{k}_n, \omega) \simeq -\left(\frac{V}{t}\right)^2 \frac{\omega^2}{\mu} \ln \left[\min \left(\frac{\omega t^2}{\Delta^2\mu}; \frac{\mu}{\omega} \right) \right]. \quad (13)$$

Again, we obtain the crossover from the cubic regime to the quadratic one. We see that close to the half-filling (at $\mu \ll t$) the energy scale of this crossover ($\Delta^2\mu/t^2$) is much less than that previously considered. It is connected with large curvature of FS near the nodal points so that the linear approximation for the spectrum breaks down much earlier. Note that this narrowing of the ω^3 region is accompanied by the strong parametric enhancement of $\text{Im}\Sigma$ by factor t^2/μ^2 .

At higher frequencies $\mu \ll \omega \ll t$ we get

$$\text{Im}\Sigma(\mathbf{k}_n, \omega) \simeq -\left(\frac{V}{t}\right)^2 \omega \ln \left[\min \left(\frac{\omega}{\mu}; \frac{t}{\omega} \right) \right] \quad (14)$$

Comparing this formula with formula found in previous subsection we see that the additional logarithmic factor exists stemming from closeness of saddle-point to the nodal one.

Note also that under $\mu \rightarrow 0$ we deal with poor-defined excitations since as it follows from our calculations $\text{Im}\Sigma \sim \omega \ln \omega$ for all ω in this case. From Eq. (14) and Kramers-Kronig relations one can find that the quasiparticles are well-defined if

$$\left[\frac{V}{t} \ln \frac{t}{\mu} \right]^2 \leq 1 \quad (15)$$

which is nothing but the criterion of absence of spin density wave. We note here that the same criterion arises in the case considered in Subsect. B, as well. The additional logarithmic enhancement stems in that case from the self-energy behavior near the “corners” of FS (cf. [8]).

IV. DISCUSSION AND CONCLUSIONS

All the results presented in previous section may be rewritten in an universal form. To do that more compactly let us introduce the following variables $a = v_g^2/(v_F \mathcal{K})$ and $b = \min(v_g, v_g/\mathcal{K})$, where $\mathcal{K}$ is the curvature of FS at the nodal point. We define also α as the angle between $\mathbf{v}_F$ and $\mathbf{v}_g$. Then we may write down the following formula:

$$\text{Im}\Sigma(\mathbf{k}_n, \omega) \simeq -\frac{V^2}{v_F^2 v_g^2 \sin^2 \alpha} \times \begin{cases} \omega^3, & \omega \ll \min(a, b); \\ \omega^2 a \ln \left[\min \left(\frac{\omega}{a}, \frac{b^2}{a\omega} \right) \right], & \min(a, b) \ll \omega \ll \max \left(b, \frac{b^2}{a} \right); \\ \omega b^2 \times \left(1 \text{ or } \ln \frac{a\omega}{b^2} \right), & \frac{b^2}{a} \ll \omega, \end{cases} \quad (16)$$

This result is valid until $v_g \ll v_F$ and $\alpha \neq 0$, as it takes place in all principal cases. The additional extrafactor $\ln(a\omega/b^2)$ at high frequencies $\omega \gg b^2/a$ appears only when the nodal point lies near the saddle one. It is easy to see that all previous results are reproduced by the following substitutions (i) $\mathcal{K} \sim 1$, $v_g \sim \Delta$, $v_F \sim \varepsilon_F$ (assuming $q_F \sim 1$) for circular FS, (ii) $\mathcal{K} \sim \mu/t$, $v_g \sim \Delta$, $v_F \sim t$ and (iii) $\mathcal{K} \sim \sqrt{t/\mu}$, $v_g \sim \Delta\sqrt{\mu/t}$, $v_F \sim \sqrt{\mu t}$ for FS of varying curvature. We mention also that the quadratic regime disappears completely when $\mathcal{K} \ll v_g/v_F$ and that the intermediate crossover scales differ principally as $\mathcal{K} < 1$ and $\mathcal{K} > 1$.

Note that we have evaluated the damping using spectrum expansion near specific points up to the second order terms in deviations from this points. This is obviously valid for Fermi surfaces smooth enough.

We undertake our consideration for circular and simplest tight-binding FSs. Meanwhile, in the adequate theory of high-temperature superconductivity the next-nearest-neighbor hopping terms in the tight-binding scheme should be taken into account. [10] It is clear from our analysis how the picture will be modified in this case. Say, for $d_{x^2-y^2}$ -wave pairing the range of cubic regime will be shrunken and a set of intermediate crossovers may appear in the range of the quadratic one. However, they will differ only by the logarithmic factor. Next, for the special choice of the parameters $2t' \approx t$ and $\mu \simeq 4t'$ the saddle-points near $(0, \pm\pi)$ and $(\pm\pi, 0)$ appear to be the “extended” ones (quasi-1D). [11] Here t' is the next-to-nearest neighbor hopping integral and μ controls the filling. This case should be treated separately.

The above consideration was made for zero temperature. It can be trivially generalized onto low finite temperatures (less than ω and minimal frequency scale in (16)). In the opposite limiting case (but not very close to T_c) Eqs. 8-14 can be rewritten by substitution T for ω . In particular, it leads to the cubic T -dependence of damping for sufficiently small temperatures: $\omega \ll T \ll \min(a, b)$ (the temperature dependence of the gap function yields only the corrections for this law). In order to apply our results to the normal state one should put simply $v_g = 0$. Note that divergence in the low-temperature asymptotics never exist, since always $T_c \gg a$. If the crossover scales $\min(a, b)$ or $\max(b, b^2/a)$ are found below T_c , T^2 or T regimes should appear in the superconducting phase. When $b^2/a \gg T_c$, the crossover from the conventional 2D FL behavior to the linear one occurs in the normal state.

Our treatment has demonstrated, in particular, that the “purely nodal” regime for damping takes place in the quite narrow frequency (temperature) range. It is restricted from above by the much less scale than the gap value as one could expect. It seems to be not a feature of damping only, so that the same characteristic scale should appear for other quantities such as density of states, nuclear relaxation rate, penetration depth, inelastic scattered neutron intensity near incommensurate and internodal vectors, *etc.* Since, on the other hand, the

impurities scattering restricts this range from below, one can estimate it as follows:

$$\Gamma_{imp} \lesssim (\omega, T) \lesssim \varepsilon_n \left[\frac{v_g}{v_F} \right]^2, \\ \varepsilon_n = \left(\frac{q_F^2}{2m} \right)_{\mathbf{q}=\mathbf{q}_n}, \quad (17)$$

where Γ_{imp} is the damping due to impurities. The estimates show that the range defined by Eq. (17) appear to be quite narrow for the high- T_c compounds. It may even disappear if Γ_{imp} will be large enough ($\sim 10\text{meV}$).

In conclusion, we consider the damping of excitations in the 2D superconductor with nodal points in the spectrum. We show that at $T = 0$ this damping demonstrates gapless cubic-in- ω low-frequency behavior at the nodal points. The results obtained for high frequencies depend on details of Fermi surface parametrization. We have analyzed two important particular FS parametrizations such as the models of almost free and tight-binding electrons. We find the frequency variation of damping and the crossover scales for these cases. For smooth enough FS they may be expressed through the model-free characteristics of the spectrum at the nodal points. We argue that the excitations are well-defined in the wide frequency range. We extend our treatment to finite temperatures and study the possibility of crossovers in temperature dependence of damping below the superconducting transition.

ACKNOWLEDGMENTS

We thank S. V. Maleyev for useful discussions and critical reading of the manuscript. The financial support from the International Science Foundation (Grant No. R3Y000) is acknowledged.

APPENDIX A:

Here we study Eq. (5) for circular FS: $\xi_{\mathbf{q}} = q^2/2m - \varepsilon_F$. We shall assume that $V_{\mathbf{k},\mathbf{q};\mathbf{p},\mathbf{l}}$ is finite in the important areas of the integrand in Eq. (5). Also, let us take for a moment the

combination of $u - v$ coefficients in Eq. (5) to be of order of unity. Then we should analyze the following expression

$$\text{Im}\Sigma \simeq V^2 \iint_D d^2\mathbf{q} d^2\mathbf{p} \delta(\varepsilon_{\mathbf{q}} + \varepsilon_{\mathbf{p}} - \omega + \varepsilon_{\mathbf{q}+\mathbf{p}-\mathbf{k}}), \quad (\text{A1})$$

where D is defined by the condition

$$\varepsilon_{\mathbf{q}} + \varepsilon_{\mathbf{p}} \leq \omega. \quad (\text{A2})$$

It is convenient to make the substitution $\hat{q} = q^2 - q_F^2$, $\hat{p} = p^2 - q_F^2$ in Eq. (A1), then $d^2\mathbf{q} = d\varphi d\hat{q}$, $d^2\mathbf{p} = d\psi d\hat{p}$, where φ and ψ are the corresponding polar angles. Taking $q_F \sim 1$, we may represent the normal-state dispersion as $\xi_{\mathbf{q}} \simeq \varepsilon_F \hat{q}$. Besides, we expand the gap functions up to the linear in angle terms $\Delta_{\mathbf{q}} \simeq \Delta\varphi$, $\Delta_{\mathbf{p}} \simeq \Delta\psi$. Using such parametrization and leaving the principal terms only, we reduce Eq. (A1) to the following one

$$\begin{aligned} \text{Im}\Sigma \simeq V^2 \int_0^{\min(1, \omega/\Delta)} \int_0^{\omega/\varepsilon_F} d\varphi d\psi \int_0^{\omega/\varepsilon_F} dq dp \delta \left(\sqrt{q^2 \varepsilon_F^2 + \Delta^2 \varphi^2} \right. \\ \left. + \sqrt{p^2 \varepsilon_F^2 + \Delta^2 \psi^2} - \omega + \varepsilon_F \varphi \psi \right), \end{aligned} \quad (\text{A3})$$

where the limits of integration stem from inequality (A2). This condition means, in particular, that at small ω momenta integrations in Eq. (A3) are restricted by the elliptic areas whose centers lie at the nodal points. It is convenient to pass therefore to the elliptic coordinates in Eq. (A3):

$$\begin{aligned} r &= \sqrt{q^2 \varepsilon_F^2 + \Delta^2 \varphi^2} & R &= \sqrt{p^2 \varepsilon_F^2 + \Delta^2 \psi^2} \\ \alpha &= \arccos q \varepsilon_F / r & \beta &= \arccos p \varepsilon_F / R \end{aligned} \quad (\text{A4})$$

Then it transforms to

$$\begin{aligned} \text{Im}\Sigma \simeq \frac{V^2 \omega^3}{\varepsilon_F^2 \Delta^2} \int_0^{\min(1, \Delta/\omega)} \int_0^1 d\alpha d\beta \int_0^1 dr dR \\ \times \delta \left(\frac{r + R - 1}{rR} + \frac{\varepsilon_F}{\Delta^2} \omega \alpha \beta \right) \end{aligned} \quad (\text{A5})$$

This expression allows us to analyze all principal limiting cases.

(1) $\omega \ll \Delta^2/\varepsilon_F$. In this case we can neglect the second term in δ -function. The integral

remained can be easily evaluated and the result is given by Eq. (9)

(2) $\Delta^2/e_F \ll \omega \ll \Delta$. Here it is convenient to rewrite δ -function in Eq. (A5) as follows

$$\frac{\Delta^2}{\omega \varepsilon_F} \delta \left(\alpha \beta + O \left(\frac{\Delta^2}{\omega \varepsilon_F} \right) \right). \quad (\text{A6})$$

One can see that the integration over one of the angles gives the logarithmic divergence which is cut on the scale $\Delta^2/\omega \varepsilon_F$.

(3) $\Delta \ll \omega \ll \varepsilon_F$. Then we should integrate the δ -function in Eq. (A5) with respect to the angles between the limits 0 and Δ/ω . After the trivial substitution $(\alpha, \beta) \mapsto (\Delta/\omega)(\alpha, \beta)$, we have

$$\frac{\omega}{\varepsilon_F} \delta \left(\alpha \beta + O \left(\frac{\omega}{\varepsilon_F} \right) \right) \quad (\text{A7})$$

Again, we have the logarithmic divergence, which is cut now on the scale ω/ε_F . The last two cases we may be combined as Eq. (10).

It can be checked out that taking the combination of $u - v$ coefficients in Eq. (5) into account does not change our estimates.

Other kinds of Fermi surface and gap functions considered in the text can be analyzed on the similar manner.

REFERENCES

- [1] See, for example: J. P. Lu, Phys. Rev. Lett. **67**, 125 (1991); N. Bulut and D. J. Scalapino, *ibid.* **68**, 706 (1992); D. S. Rokhsar, *ibid.* **70**, 493 (1993); C.-R. Hu, *ibid.* **72**, 1526 (1994); J. P. Lu, J. Mod. Phys. Lett. B **6**, 547 (1992); P. Monthoux *et al.*, Phys. Rev. B **47**, 6069 (1993); D. Telen, D. Pines, and J. P. Lu, *ibid.* 9151 (1993); T. P. Devereaux and P. Fulde, *ibid.* 14638 (1993); P. J. Hirshfeld and N. Goldenfeld, *ibid.* **48**, 4219 (1993); H. Won and K. Maki, *ibid.* **49**, 15305 (1994); S. V. Maleyev, A. G. Yashenkin, and D. N. Aristov, *ibid.* **50**, 13825 (1994); A. J. Leggett, Physica B **199 & 200**, 291 (1994); P. Monthoux and D. J. Scalapino, *ibid.* 294 (1994); D. Z. Liu, K. Levin, and J. Maly, preprint; A. M. Tikofsky and D. B. Bailey, preprint.
- [2] J. M. Valles *et al.*, Phys. Rev. B **44**, 11986 (1991); D. A. Bonn *et al.*, *ibid.* **47**, 11314 (1993); D. A. Brawner and H. R. Ott, *ibid.* **50**, 6530 (1994); W. N. Hardy *et al.*, Phys. Rev. Lett. **70**, 3999 (1993); D. A. Wollman *et al.*, *ibid.* **71**, 2134 (1993); P. Chaudhary and S.-Y. Lin, *ibid.* **72**, 1084 (1994); C. C. Tsuei *et al.*, *ibid.* **73**, 593 (1994); K. Moler *et al.*, *ibid.* 2744 (1994).
- [3] P. A. Lee, Phys. Rev. Lett. **71**, 1887 (1993); A. A. Nersesyan, A. M. Tsvelik, and F. Wenger, *ibid.* **72**, 2628 (1994); A. V. Balatsky, A. Rosengren, and B. L. Altshuler, *ibid.* **73**, 720 (1994).
- [4] G. Baskaran, Z. Zou and P. W. Anderson, Solid State Commun. **63**, 973 (1987); L. B. Ioffe and A. I. Larkin, Phys. Rev. B **39**, 8988 (1989).
- [5] G. Kotliar, Phys. Rev. B **37**, 3664 (1988).
- [6] N. E. Bickers, D. J. Scalapino, and R. T. Scalettar, Int. J. Mod. Phys. B **1**, 687 (1987); P. A. Lee and N. Read, Phys. Rev. Lett. **58** 2691 (1987); P. Monthoux, A. V. Balatsky, and D. Pines, *ibid.* **67**, 3448 (1991).
- [7] A. V. Chaplik, Zh. Eksp. Teor. Fiz. **60**, 1845 (1971) [Sov. Phys. JETP **33**, 997 (1971)];

- C. Hodges, H. Smith, and J. Wilkins, Phys. Rev. B **4**, 302 (1971).
- [8] D. N. Aristov, S. V. Maleyev, and A. G. Yashenkin, Phys. Rev. B **48**, 3527 (1993).
- [9] See, for example: F. Wenger and S. Östlund, Phys. Rev. B **47**, 5977 (1993).
- [10] Q. Si *et al.*, Phys. Rev. B **47**, 9055 (1993).
- [11] See, for example: A. A. Abrikosov, J. C. Campuzano and K. Gofron, Physica C **214**, 73 (1993).

FIGURES

FIG. 1. The disposition of the momenta which give the principal contribution to $\text{Im}\Sigma$ at $\omega \rightarrow 0$. The solid lines denote the Fermi surface and the dashed lines correspond to the nodes of the gap function.

FIG. 2. The disposition of the momenta which give the principal contribution to $\text{Im}\Sigma$ at $\omega \gg \mu$ in the case of nested FS. The crosses label the saddle-points. The nodal points are located (a) far away and (b) near the saddle ones.





